

SCIENTIFIC REPORTS

OPEN

Fine particulate matter (PM_{2.5}) in China at a city level

Yan-Lin Zhang & Fang Cao

Received: 24 June 2015

Accepted: 02 September 2015

Published: 15 October 2015

This study presents one of the first long term datasets including a statistical summary of PM_{2.5} concentrations obtained from one-year monitoring in 190 cities in China. We found only 25 out of 190 cities could meet the National Ambient Air Quality Standards of China, and the population-weighted mean of PM_{2.5} in Chinese cities are 61 µg/m³, ~3 times as high as global population-weighted mean, highlighting a high health risk. PM_{2.5} concentrations are generally higher in north than in south regions due to relative large PM emissions and unfavorable meteorological conditions for pollution dispersion. A remarkable seasonal variability of PM_{2.5} is observed with the highest during the winter and the lowest during the summer. Due to the enhanced contributions from dust particles and open biomass burning, high PM_{2.5} abundances are also found in the spring (in Northwest and West Central China) and autumn (in East China), respectively. In addition, we found the lowest and highest PM_{2.5} often occurs in the afternoon and evening hours, respectively, associated with daily variation of the boundary layer depth and anthropogenic emissions. The diurnal distribution of the PM_{2.5}-to-CO ratio consistently displays a pronounced peak during the afternoon periods, reflecting a significant contribution of secondary PM formation.

Due to a rapid economic development, industrial expansion and urbanization during the last few decades in China, increasingly occurrence of haze or smog episodes characterized by the high fine particulate matter (i.e. with aerodynamic diameters not larger than 2.5 µm, or PM_{2.5}) levels and reduced visibility has been reported in national-scale China especially in the most developed and high-populated city clusters such as the Yangtze River Delta (YRD), the Pearl River Delta (PRD) and the Beijing-Tianjin-Hebei (BTH) regions^{1–7}. Urban PM_{2.5} originates mainly from sources such as traffic-related emissions, road/soil dust, biomass burning, and agriculture activities as well as regional transported aerosols, but it still remains much challenging to quantify contributions from each source and understand PM formation mechanism^{2,7–11}. According to a report from Asian-development Bank, only <1% of 500 largest cities of China could meet the air quality guideline (10 µg/m³ for annual mean and 25 µg/m³ for 24-hour mean) suggested by the World Health Organization and several cities are ranked among the most polluted cities in the world¹².

High occurrence of extreme haze episodes in recent years not only leads to a global concern due to its adverse health effects^{13,14}, but also triggers the Chinese government to tackle the serious air quality problem especially PM_{2.5} pollution. From February 29th, 2012, the Chinese Ministry of Environmental Protection (MEP) published the third revision of the “the national ambient air quality standards” (NAAQS) (GB3095-2012), in which PM_{2.5} is included into the NAAQS for the first time. PM_{2.5} monitoring has not yet been introduced in the national network in China before the new NAAQS, although PM_{2.5} levels have been reported in research studies in some developed cities such as Beijing, Shanghai, Chongqing and Guangzhou^{7,8,15–18}. Due to the lack of large-scale PM_{2.5} monitoring, the spatial distribution of urban PM_{2.5} in China is often retrieved by the satellite remote sensing¹⁹. Large scale real-time PM_{2.5} monitoring is very necessary to study spatial and temporal (i.e. seasonal and diurnal) variation of urban PM_{2.5} in China. Wang *et al.* (2014) reported a study on the spatial and temporal variations of the 6 criteria pollutants in 31 provincial capital cities in China during 2013–2014²⁰. The study reveals

Yale-NUIST Center on Atmospheric Environment, Nanjing University of Information Science and Technology, Nanjing 10044, China. Correspondence and requests for materials should be addressed to Y.L.Z. (email: dryanlinzhang@gmail.com)

PM_{2.5}, PM₁₀, CO and SO₂ concentrations are higher in the North region than those in the West and the South-East regions, although high pollution events are also frequently observed during the autumn for the South-East region and during the spring for the West region. However, little has been reported regarding diurnal patterns in aerosol concentrations in different seasons due to the lack of real-time monitoring. In addition, a higher spatial-temporal resolution is also needed to obtain a better understanding of air quality problem at a city-level.

This paper presents one of the first long term data sets including a statistical summary of PM_{2.5} concentrations measured during one-year continuous monitoring in 190 priority pollution monitoring cities in China. A national air quality monitoring network with nearly 950 monitoring stations in 190 Chinese cities is put into operation by the end of 2013, which release real-time monitoring data on air quality such as PM_{2.5}, PM₁₀, SO₂, NO₂, CO, O₃ to the public. In this study, we will present an overview of spatial and seasonal distribution of PM_{2.5} in 190 cities of China. In addition, diurnal variation of PM_{2.5} in the most populated and developed cities of the PRD, YRD and BTH regions will be discussed. Finally, the population-weighted mean of PM_{2.5} and cumulative population distribution in Chinese cities will be estimated. By studying this extensive dataset at a national-scale level, we could improve our understanding of urban PM_{2.5} pollution at a fine spatiotemporal resolution in China. The obtained knowledge in this study is therefore very useful for the implementation of national/regional air pollution reduction policy in order to meet Chinese new NAAQS.

Results Overview and spatial distributions of PM_{2.5} pollution in China

Figure 1a shows spatial distributions of annual mean of PM_{2.5} concentrations of 190 cities in China. The new NAAQS of China sets PM_{2.5} concentration limits for both the 24-hour average (35 μg/m³ and 75 μg/m³ for Grade I and II, respectively) and the annual mean value (15 μg/m³ and 35 μg/m³ for Category I and II zones, respectively)²¹. The annual 24-h PM_{2.5} ranges from 18 μg/m³ (in Sanya) to 116 μg/m³ (in Baoding) with average of 57 ± 18 μg/m³, severely exceeding the NAAQS of China and other standards recommended by international organizations and other countries (10–35 μg/m³)^{21–23}. According to the new NAAQS (Grade II), as many as 165 cities cannot meet the standards, accounting for 87% of the total number of cities. PM_{2.5} concentrations are generally higher in the cities located in the North regions than those observed in the South regions. PM_{2.5} concentrations also tend to be lower in the coastal regions than in the inland regions. As can be seen in Fig. 1a, the highest annual average concentration is observed in the Beijing-Tianjin-Hebei (BTH) region including two megacities of Beijing and Tianjin and Hebei Province, which has the highest density of coal consumption and heavy industries (e.g. iron, steel and cement) in China. It is important to note that half of the 10 worst haze affected cities are in Beijing-Tianjin-Hebei, namely Baoding, Xingtai, Shijiazhuang, Handan and Hengshui (see Table S1). Several studies have revealed that the enhanced PM pollution in BTH is not only due to the primary emissions from local sources such as industrial and domestic and agricultural sources but also due to the regional transported contribution (e.g. from nearby Shandong and Henan Provinces) and secondary production^{8,9,17,18,24,25}. Furthermore, the climate of BTH is characterized by stagnant weather with weak wind and relatively low boundary layer height, leading to the favorable atmosphere conditions for accumulation, formation and processing of aerosols²⁶. The lowest PM_{2.5} concentration is observed in Hainan Province, the southernmost island of China, which is due to the less anthropogenic emissions and favorable meteorological condition for atmospheric dispersion and dilution. Due to the less coal-based industries and good dispersion weather conditions, the PM_{2.5} concentrations in PRD (i.e. cities in Guangdong province, see Table S1) are generally lower than those in the other two largest city clusters in China such as BTH and YRD (cities in Jiangsu and Zhejiang provinces as well as Shanghai, see Table S1), although 7 out of 21 cities in PRD still could not meet the new NAAQS of China.

Seasonal variation of PM_{2.5} pollution in China

Generally, PM_{2.5} concentrations show a remarkable seasonal variability with the highest during the winter and the lowest during the summer (Fig. 1). The wintertime maximum is associated with enhanced anthropogenic emissions from fossil fuel combustion and biomass burning and unfavorable meteorological conditions (i.e. more frequent occurrences of stagnant weather and temperature inversion during the cold periods) for pollution dispersion. In addition to accumulation of primary emissions, new particle formation and secondary production of both the inorganic aerosols and organic matters could further enhance fine PM abundance^{8,9}. Actually, the PM_{2.5} to PM₁₀ ratio is slightly higher during the winter than during the other seasons (Fig. 2 and S1), suggesting importance of combustion sources and secondary formation of fine particles. However, the seasonality displays a spatial variability. For Northwest and West Central China, the most polluted season appears in spring but not in winter, associated with increased contribution from dust particles in this desert-like region. PM_{2.5}/PM₁₀ ratio (i.e.0.21) during the spring in Korla (41.7N, 86.1E), one of most polluted city in West China, are much lower than all other studied cities in China (i.e. with a mean of 0.56 ± 0.10). The decrease of PM_{2.5} to PM₁₀ ratios are also observed in many cities in West Central and Northwest China (Fig. 2). The higher abundance of the coarse particles (PM_{2.5–10}) in PM₁₀ indicates a significant contribution from local dust emission and regional dust transport, as the other coarse-mode particle such as biogenic-derived large particles (e.g. plant debris) could only have a minor if not a negligible contribution with its cold desert climate and low vegetation coverage. Actually, the dust storm is most frequently occurred in the west regions in Northern China




Figure 1. The averaged $PM_{2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) of the 190 cities of China, during the year of 2014/2015 (a) and during the spring (b), summer (c), autumn (d), and winter (e). The maps were drawn by the software of Igor Pro, <http://www.wavemetrics.com/>.

in spring from recent observations where desert, semi-desert or grassland is distributed²⁷. This is also evident by seasonal variation of $PM_{2.5}/PM_{10}$ ratios (Fig. 2 and S2), suggesting significant importance of dust particles to $PM_{2.5}$ in west regions of China. The substantial fraction of PM_{10} mass in the $PM_{2.5-10}$ size range also suggests that the current $PM_{2.5}$ control strategies (i.e. reduce fossil/non-fossil combustion derived VOCs and PM emissions) will only partly reduce the PM_{10} pollution in the west part of China. $PM_{2.5}$ is much decreased during summer associated with the reduced anthropogenic emissions such as fossil fuel and biomass burning for domestic heating. Further, large wet depositions of aerosols, clean air mass from ocean during the Asian summer monsoon and enhanced convection within a higher atmospheric mixing layer could lead to a strong dilution effect of aerosols in Eastern China²⁸. As a result, the $PM_{2.5}$ minima are observed in summer. Interestingly, relatively high $PM_{2.5}$ levels are also found during the autumn over Eastern China, which is likely due to enhanced open biomass burning during the agricultural harvest season. The fire counts map derived from MODIS on the NASA satellite (Figure S2) shows that during the October, intensive open biomass burning (i.e. agricultural waste burning) events occur over Northeast and South China. It should be also noted that $PM_{2.5}$ concentrations during the




Figure 2. The averaged $PM_{2.5}/PM_{10}$ ratio of the 190 cities of China, during the year of 2014/2015 (a) and during the spring (b), summer (c), autumn (d), and winter (e). The maps were drawn by the software of Igor Pro, <http://www.wavemetrics.com/>.

autumn in the cities of the PRD and Northeast regions are higher than those found during the spring. In few cases (9 out of 190 cities), $PM_{2.5}$ is even higher during the autumn than during the winter. The importance of biomass burning contributions to $PM_{2.5}$ during autumn over PRD and Northeast China has also been underscored by previous studies^{29,30}. For example, Wang *et al.* (2007) reported up to 19% of total $PM_{2.5}$ is from biomass burning emissions in Guangzhou, the largest city in the PRD region³¹.

Diurnal variation of $PM_{2.5}$

In this study, hourly data were used to examine diurnal variability in $PM_{2.5}$ as well as the other major air pollutants. This provides importance information in identifying the potential emission sources and the time when the pollution level exceeds the standards. Figure 3 illustrates the diurnal variation of hourly $PM_{2.5}$ concentrations in each season in Beijing, Shanghai and Guangzhou, the largest megacities in BTH, YRD and PRD in different climatic zones of China, respectively.




Figure 3. Diurnal variations of hourly PM_{2.5} concentrations in Beijing (a), Shanghai (b) and Guangzhou (c).

In Beijing, PM_{2.5} shows significantly higher concentrations and pronounced diurnal variations during the autumn and winter than during the spring and summer (Fig. 3a). The PM_{2.5} concentration during the autumn and winter is higher during the nighttime than the daytime, which can be explained by the enhanced emission for heating and relatively low the boundary layer (Fig. 4). The peak PM_{2.5} concentration at night is ~2 times higher than that of valley in the afternoon. The lowest concentrations are observed in afternoon hours when the boundary layer becomes larger and wind speed increases (data not shown). After 16:00, PM_{2.5} concentrations start to increase because of the quickly decreasing boundary layer heights (Fig. 4) and increasing vehicle emissions as evident by increased NO₂ emissions (see Fig. 5). Moreover, PM pollution emitted from diesel truck traffic which is allowed only during nighttime additionally increase PM burden because emission factors of heavy-duty vehicles are 6 times than those from light-duty vehicles³². It should be noted that such a traffic restriction has been applied in many Chinese megacities which may affect diurnal pattern of PM_{2.5} and its chemical compositions. PM_{2.5} levels in afternoon periods during winter are on average even lower than those observed during the other seasons. Such a trend is not observed for other major air pollutants such as NO₂ and SO₂ (Figs 5 and 6). This indicates that the PM_{2.5} levels in Beijing are not only driven by primary emissions but are also affected by other factors such as meteorological conditions and secondary PM production. To further discuss secondary production of PM_{2.5}, CO is used to normalize PM_{2.5} concentrations to exclude the influence of primary combustion emissions and meteorological factors³³. Figure 7 shows diurnal variation of hourly ratios of PM_{2.5} to CO (i.e. an excellent tracer for primary combustion source). A pronounced peak of the PM_{2.5}/CO ratio occurs during the afternoon (i.e. around 16:00) for all the seasons although the ratio is apparently lower during the winter than during the other seasons. This indicates secondary formations




Figure 4. Daily evolution of average planetary boundary layer height in Beijing (a), Shanghai (b) and Guangzhou (c).

process also plays an important role in PM concentrations, especially in the afternoon when the photochemical activities are relatively strong. However, a more detailed explanation is subject to further studies including more comprehensive observations of chemical compositions of $PM_{2.5}$ and its precursors as well as meteorology. In spring and summer, the moderate $PM_{2.5}$ peak appears in the morning (i.e. 8:00 am to 11:00 am), in accordance with high NO_2 emissions from transportation during rush hours.

As shown in Fig. 3b, a unique diurnal pattern is observed in Shanghai. $PM_{2.5}$ concentration during the winter is higher than those found in the other seasonal throughout the day, which could be explained by the enhanced anthropogenic emissions. In addition, the heavily polluted air mass is often transported from Northern China to the YRD region under East Asian winter monsoon climate (Figure S3), affecting the regional air quality in YRD³⁴. The enhancement of SO_2 concentration in the winter is comparably higher than that of NO_2 (Figs 5 and 6), suggesting the importance of coal combustion emissions (i.e. power plant). There are often two moderate peaks of $PM_{2.5}$ concentrations: one in the morning between 7 and 10 am, and another in the evening between 7 and 10 pm. A similar morning time peak is also found for both SO_2 (i.e. a tracer for coal combustion) and NO_2 (i.e. a tracer for vehicle emissions), suggesting a significant contribution of fossil emissions (e.g. from power plant and vehicle exhaust) to daytime $PM_{2.5}$. Concentrations decrease from mid-morning to later afternoon/early evening due to a combination of the increasing boundary layer depth and reduced anthropogenic emissions. The afternoon increased $PM_{2.5}$ is consistent with observation of NO_2 , suggesting traffic-related emissions may play a more important role in the $PM_{2.5}$ variation from the afternoon to the evening hours. The diurnal distributions of $PM_{2.5}/CO$




Figure 5. Diurnal variations of hourly NO₂ concentrations in Beijing (a), Shanghai (b) and Guangzhou (c).

show lowest levels during the evening and morning hours suggesting a predominance of combustions in the rush (traffic) hours.

Similar to Beijing, a strong daily variation of PM_{2.5} in Guangzhou is found during the autumn and winter, which is characterized by the minima in the late afternoon and maxima in the night. During the autumn and winter, an increase in PM_{2.5} is not observed during the morning periods although both the SO₂ and NO₂ are increased due to the increased anthropogenic emissions. Alternatively, this decreasing may be the result of increasing boundary layer depth. In these two seasons, the PM_{2.5} levels start to increase from the later afternoon, which could be explained by the increased motor vehicle emissions as NO₂ is also dramatically increased during the morning time. It is interesting to note that the daily variation pattern of SO₂ (and NO₂) is very similar in different seasons, although PM_{2.5} displays a remarkably different diurnal cycle during the autumn and winter compared to during the spring and summer. The results indicate that PM formation process is very complex and is not only determined by the emission strength but is also driven by the other influencing factors such as the meteorological conditions and atmospheric oxidant capacity. During the spring and summer, a moderate PM_{2.5} peak often occurs in the early afternoon, typically earlier than peak hours during the autumn and winter. This phenomenon (i.e. a shift of peak time in the afternoon) is likely due to the longer and stronger solar irradiation, increasing O₃ concentrations and higher temperatures, leading to the enhanced photochemical formation of secondary aerosol particles, being important constituents of PM_{2.5}. Secondary organic aerosols formation from biogenic volatile organic compounds during spring and summer may also increase PM_{2.5} levels because the Guangzhou is characterized by the sub-tropical climate with the annual mean temperature of around 25°C and high evergreen vegetation coverage^{35,36}. The PM_{2.5} concentration keeps in high levels during the




Figure 6. Diurnal variations of hourly SO₂ concentrations in Beijing (a), Shanghai (b) and Guangzhou (c).

nighttime (i.e. ~19:00 to midnight) in all the seasons, associated with primary automobiles emissions and the subsequent accumulation and secondary PM production. The PM_{2.5}/CO ratio show a moderate daily variability with the highest value in the afternoon time during the spring and summer, because of the longer and stronger solar irradiation and increasing O₃ concentrations as discussed before. However, no significant diurnal pattern is observed for the other two seasons. It should be also noted that the PM_{2.5}/CO ratio is generally lower than those in Beijing and Shanghai, indicating a more important contribution of combustion emissions (i.e., from gasoline vehicles and biofuel) to PM_{2.5} concentration in Guangzhou and/or higher contribution of coal combustions in Beijing and Shanghai.

Discussion

This study analyzes PM concentration data collected from the newest air quality monitoring network of the Ministry of Environmental Protection of China in 190 major cities during April 2014 to April 2015. The annual averaged concentration of PM_{2.5} is $57 \pm 18 \mu\text{g}/\text{m}^3$ (ranging from 16 to $119 \mu\text{g}/\text{m}^3$), severely exceeding the new NAAQS ($35 \mu\text{g}/\text{m}^3$) of China and World Health Organization air quality guideline ($10 \mu\text{g}/\text{m}^3$). PM_{2.5} concentrations are generally higher in the cities located in the north region than those observed in the south regions, and are lower in the coastal regions than in the inland regions. This suggests that different control strategies should be targeted in different regions of China in terms of specific feature of local/regional emissions and meteorology. Generally, PM_{2.5} concentrations show a remarkable seasonal variability with the highest during the winter and the lowest during the summer. The winter maximum PM_{2.5} level is due to the increasing anthropogenic activities such as fossil-fuel and




Figure 7. Diurnal variations of hourly $PM_{2.5}$ to CO ratios ($PM_{2.5}/CO$) concentrations in Beijing (a), Shanghai (b) and Guangzhou (c).

biomass burning for heating in the cold season. Furthermore, more frequent occurrences of stagnant weather and temperature inversion during the cold period may facilitate the accumulation of air pollutants. Alternatively, more frequent occurrences of precipitation and mixing of transported clean air mass result in a strong deposition, dispersion and dilution of $PM_{2.5}$ in summer. It should be also noted the most polluted season appears in spring but not in winter for some cities located in Northwest and West Central China, associated to a higher contribution from dust particles in this desert-like region. Moderately high $PM_{2.5}$ is also observed during the autumn due to the enhanced open biomass burning emissions in the harvest season. The diurnal pattern of $PM_{2.5}$ concentration as well as other major air pollutants such as NO_2 , SO_2 and CO is also characterized. In Beijing and Guangzhou, a pronounced daily variation of $PM_{2.5}$ concentration is found during the autumn and winter, with the lowest levels in the afternoon hours, apparently due to the deeper boundary layer. The $PM_{2.5}$ keeps in high abundance at the evening hours in the cold seasons because of increasing emissions for heating and stagnant atmospheric conditions. Finally, CO is used to normalize $PM_{2.5}$ concentrations to exclude the influence of primary combustion emissions and meteorological factors. The diurnal distribution of the $PM_{2.5}$ to CO ratio (i.e. $PM_{2.5}/CO$) consistently displays a remarkable peak during the afternoon periods, reflecting a significant contribution of secondary particle formation due to the relatively stronger solar irradiation and higher temperature. Our results demonstrate that the spatial-temporal distribution of $PM_{2.5}$ in China is determined by complex factors such as local/regional-transported emissions, meteorological conditions and atmospheric oxidant capacity. In order to obtain a more comprehensive picture of source and formation




Figure 8. Cumulative distribution of annual mean $PM_{2.5}$ estimated from ground measurements in 190 Chinese cities. The results from the Beijing-Tianjin-Hebei (BTH), Yangtze River Delta (YRD), and Pearl River Delta (PRD) regions are also displayed.

mechanism of fine particulate matter pollution over national wide China, the real-time chemical composition of $PM_{2.5}$ and the major air pollutants (e.g. NO_x , SO_2 , O_3 , CO and volatile organic compounds) as well as meteorological conditions should be simultaneously investigated in a fine spatial-temporal resolution in future.

A major challenge for assessments of $PM_{2.5}$ -associated health impacts often lies in the lack of ground-based monitoring networks and representative exposure estimates¹⁹. This paper provides one of first datasets including $PM_{2.5}$ levels in 190 Chinese cities from ~950 monitoring sites, which could be used for assessments of air pollution health impacts. Based on annual 24-hour $PM_{2.5}$ and population data in each studied city, the population-weighted mean of $PM_{2.5}$ in Chinese cities are $61 \mu\text{g}/\text{m}^3$, approximately 3 times as high as global population-weighted mean ($20 \mu\text{g}/\text{m}^3$)¹⁹. Figure 8 shows the cumulative population distribution in Chinese cities as well as the most three developed regions such as the BTH, YRD and PRD regions. Overall, only 9% of the population live in cities with annual $PM_{2.5}$ mean smaller than the new NAAQS in China ($35 \mu\text{g}/\text{m}^3$). It should be noted that all the population live in cities where the WHO Air Quality Guideline ($10 \mu\text{g}/\text{m}^3$) is exceeded. All studied regions show a nonlinear relationship of $PM_{2.5}$ concentrations with population, which is consistent with the previous findings conducted in other places around the world¹⁹. The cities in the BTH region have the highest levels of $PM_{2.5}$ concentrations, with 70% of the regional population exceeding $70 \mu\text{g}/\text{m}^3$ (twice as high as the NAAQS of China), whereas in the relatively clean region (i.e., PRD) there is still 70% of population residing in the cities exceeding the NAAQS of China (i.e., $35 \mu\text{g}/\text{m}^3$). Hoek *et al.* (2013) estimated that an increase of $10 \mu\text{g}/\text{m}^3$ in long-term $PM_{2.5}$ exposure increases a 6.2% (95% confidence interval 4.1–8.4%) of all-cause mortality based on meta-analysis of cohort studies³⁷. Taken together, this study highlights the significantly high health risk from fine particulate matter pollutions and difficulties in achieving air quality targets in China. Therefore, China needs much more effective policies and research that could mitigate complex $PM_{2.5}$ pollutions, reduce urban population exposures, and achieve sustainable development. Future epidemiologic studies are urgently required to provide direct estimation of relationship of $PM_{2.5}$ and mortality such as chronic cardiovascular and respiratory diseases in China and other polluted countries/regions.

Methods

The national air quality monitoring network is continuously operated and maintained by Department of the Environment for each city. The network comprises of 496 stations in 74 cities since 2012, which is now extended to 946 monitoring stations in 190 cities (Table S1 and S2) from 2013. At each monitoring site, the real-time mass concentrations of $PM_{2.5}$ and PM_{10} are measured using the micro oscillating balance method and/or the β absorption method from commercial instruments. Gas pollutants such as SO_2 , NO_2 , CO are measured using the ultraviolet fluorescence method, the chemiluminescence method and the gas filter correlation infrared absorption (or the non-dispersive infrared absorption method), respectively, by a set of commercial instruments. The instrumental operation, maintenance, data assurance and quality control are properly conducted according to the most revisions of China Environmental Protection Standards such as “HJ 193–2013”³⁸ and “HJ 655–2013”³⁹. The real-time hourly $PM_{2.5}$ concentrations data as well as other major air pollutant (i.e. SO_2 , NO_2 , $PM_{2.5}$, PM_{10} , CO, and O_3) are continuously recorded by the MEP in China and are publicly accessible⁴⁰. The data present in this study was obtained from the website during the period from 12-Apr-2014 to 11-Apr-2015. The hourly and 24-h (daily) concentrations of all the air pollutants were the averages of the hourly data from all monitoring sites in the city. Time-series (3-hour resolution) of planetary boundary layer depths were obtained from the U.S. National Oceanic and Atmospheric Administration (NOAA) READY archived meteorological GDAS data ($1^\circ \times 1^\circ$) based on Coordinated Universal Time (UTC). All UTC values are

converted to local time (Beijing time, UTC + 8). Statistical analyses are carried out using t-test (Excel 2010) when comparisons are made.

References

- Zhang, R., Li, Q. & Zhang, R. Meteorological conditions for the persistent severe fog and haze event over eastern China in January 2013. *SCIENCE CHINA Earth Sciences* **57**, 26–35 (2014).
- Zhang, Y. L. *et al.* Fossil vs. non-fossil sources of fine carbonaceous aerosols in four Chinese cities during the extreme winter haze episode of 2013. *Atmos. Chem. Phys.* **15**, 1299–1312 (2015).
- Zhang, Q., Streets, D. G., He, K. & Klimont, Z. Major components of China's anthropogenic primary particulate emissions. *Environ. Res. Lett.* **2**, 045027 (2007).
- Wang, G. H., Kawamura, K., Lee, S., Ho, K. F. & Cao, J. J. Molecular, seasonal, and spatial distributions of organic aerosols from fourteen Chinese cities. *Environ. Sci. Technol.* **40**, 4619–4625 (2006).
- Cao, J. J. *et al.* Spatial and seasonal distributions of carbonaceous aerosols over China. *J. Geophys. Res.* **112**, D22S11 (2007).
- Cao, J. J. *et al.* Winter and summer PM_{2.5} chemical compositions in fourteen Chinese cities. *J. Air Waste Manage. Assoc.* **62**, 1214–1226 (2012).
- Chan, C. K. & Yao, X. Air pollution in mega cities in China. *Atmos. Environ.* **42**, 1–42 (2008).
- Huang, R. J. *et al.* High secondary aerosol contribution to particulate pollution during haze events in China. *Nature* **514**, 218–222 (2014).
- Guo, S. *et al.* Elucidating severe urban haze formation in China. *Proc. Nat. Acad. Sci. U.S.A.* **111**, 17373–17378 (2014).
- Zhang, Y. L. *et al.* Source Apportionment of Elemental Carbon in Beijing, China: Insights from Radiocarbon and Organic Marker Measurements. *Environ. Sci. Technol.* **49**, 8408–8415 (2015).
- He, H. *et al.* Mineral dust and NO_x promote the conversion of SO₂ to sulfate in heavy pollution days. *Sci Rep* **4**, 4172 (2014).
- Zhang, Q. & Crooks, R. *Toward an environmentally sustainable future: Country environmental analysis of the People's Republic of China.* (Asian Development Bank, 2012).
- Pope III, C. A. & Dockery, D. W. Health effects of fine particulate air pollution: Lines that connect. *J. Air Waste Manage. Assoc.* **56**, 709–742 (2006).
- Pope, C. A. & Dockery, D. W. Air pollution and life expectancy in China and beyond. *Proc. Nat. Acad. Sci. USA* **110**, 12861–12862 (2013).
- Yang, F. *et al.* Characteristics of PM_{2.5} speciation in representative megacities and across China. *Atmos. Chem. Phys.* **11**, 5207–5219 (2011).
- Yang, F. *et al.* One-year record of organic and elemental carbon in fine particles in downtown Beijing and Shanghai. *Atmos. Chem. Phys.* **5**, 1449–1457 (2005).
- Zhao, P. S. *et al.* Characteristics of concentrations and chemical compositions for PM_{2.5} in the region of Beijing, Tianjin, and Hebei, China. *Atmos. Chem. Phys.* **13**, 4631–4644 (2013).
- Zhang, J. K. *et al.* Characterization of submicron aerosols during a month of serious pollution in Beijing, 2013. *Atmos. Chem. Phys.* **14**, 2887–2903 (2014).
- Donkelaar, A. v. *et al.* Global estimates of ambient fine particulate matter concentrations from satellite-based aerosol optical depth: development and application. *Environ. Health Perspect.* **118**, 847 (2010).
- Wang, Y., Ying, Q., Hu, J. & Zhang, H. Spatial and temporal variations of six criteria air pollutants in 31 provincial capital cities in China during 2013–2014. *Environ. Int.* **73**, 413–422 (2014).
- Chinese Ministry of Environmental Protection (MEP). *Ambient Air Quality Standards (GB 3095-2012).* (2012) Available at <http://kjs.mep.gov.cn/hjbhbz/bzwb/dqjhbd/dqjhzbz/201203/W020120410330232398521.pdf>. (Accessed: 5th August 2015).
- EPA, U. S. 40 CFR Parts 50, 51, 52, 53, and 58-National Ambient Air Quality Standards for Particulate Matter: Final Rule. *Fed. Regist.* **78**, 3086–3286 (2013).
- WHO. *Air Quality Guidelines: Global Update 2005: Particulate Matter, Ozone, Nitrogen Dioxide and Sulfur Dioxide.* (World Health Organization, 2006).
- Sun, Y. L. *et al.* Investigation of the sources and evolution processes of severe haze pollution in Beijing in January 2013. *J. Geophys. Res.* **119**, 4380–4398 (2014).
- Wang, L. T. *et al.* The 2013 severe haze over southern Hebei, China: model evaluation, source apportionment, and policy implications. *Atmos. Chem. Phys.* **14**, 3151–3173 (2014).
- Chen, Z. H. *et al.* Relationship between atmospheric pollution processes and synoptic pressure patterns in northern China. *Atmos. Environ.* **42**, 6078–6087 (2008).
- Zou, X. K. & Zhai, P. M. Relationship between vegetation coverage and spring dust storms over northern China. *J. Geophys. Res.* **109**, D03104 (2004).
- Yoo, J.-M. *et al.* New indices for wet scavenging of air pollutants (O₃, CO, NO₂, SO₂, and PM₁₀) by summertime rain. *Atmos. Environ.* **82**, 226–237 (2014).
- Zhang, Z. S. *et al.* Characteristics and applications of size-segregated biomass burning tracers in China's Pearl River Delta region. *Atmos. Environ.* **102**, 290–301 (2015).
- Zhang, Y. L. & Cao, F. Is it time to tackle PM(2.5) air pollutions in China from biomass-burning emissions? *Environ. Pollut.* **202**, 217–219 (2015).
- Wang, Q. Q. *et al.* Impact of biomass burning on urban air quality estimated by organic tracers: Guangzhou and Beijing as cases. *Atmos. Environ.* **41**, 8380–8390 (2007).
- Westerdahl, D., Wang, X., Pan, X. C. & Zhang, K. M. Characterization of on-road vehicle emission factors and microenvironmental air quality in Beijing, China. *Atmos. Environ.* **43**, 697–705 (2009).
- de Gouw, J. A. *et al.* Emission and chemistry of organic carbon in the gas and aerosol phase at a sub-urban site near Mexico City in March 2006 during the MILAGRO study. *Atmos. Chem. Phys.* **9**, 3425–3442 (2009).
- Li, M. M. *et al.* Analysis of the transport pathways and potential sources of PM₁₀ in Shanghai based on three methods. *Sci. Tot. Environ.* **414**, 525–534 (2012).
- Wang, X. M. *et al.* Spatiotemporal variability of biogenic terpenoid emissions in Pearl River Delta, China, with high-resolution land-cover and meteorological data. *Tellus Series B-Chemical and Physical Meteorology* **63**, 241–254 (2011).
- Ding, X. *et al.* Tracer-based estimation of secondary organic carbon in the Pearl River Delta, south China. *J. Geophys. Res.* **117**, D05313 (2012).
- Hoek, G. *et al.* Long-term air pollution exposure and cardio- respiratory mortality: a review. *Environmental Health* **12**, doi: 10.1186/1476-069x-12-43 (2013).
- Chinese Ministry of Environmental Protection (MEP)., Technical Specifications for Installation and Acceptance of Ambient air Quality Continuous Automated Monitoring System for SO₂, NO₂, O₃ and CO. (2013) Available at <http://www.es.org.cn/download/2013/7-12/2627-1.pdf>. (Accessed: 5th August 2015).

39. Chinese Ministry of Environmental Protection (MEP). Technical Specifications for Installation and Acceptance of Ambient Air Quality Continuous Automated Monitoring System for PM₁₀ and PM_{2.5}. (20130) Available at <http://kjs.mep.gov.cn/hjbhbz/bzwb/dqjhbjh/jcgfffbz/201308/W020130802492823718666.pdf>. (Accessed: 5th August 2015).
40. China's National Environmental Monitoring Center. *The real-time air quality monitoring data*. (2013) Available at <http://113.108.142.147:20035/emcpublish/China>, (Accessed: 5th August 2015).

Acknowledgment

We acknowledge the financial support from the Natural Science Foundation for Young Scientists of Jiangsu Province, China (BK20150895) and the Startup Foundation for Introducing Talent of NUIST (No. 2015r023 and 2015r019). We also acknowledge the use of fire/hotspot data produced by the University of Maryland and provided by NASA FIRMS operated by NASA/GSFC/ESDIS with funding provided by NASA/HQ (available on-line <https://earthdata.nasa.gov/active-fire-data#tab-content-6>). The support of Xiao-Yan Liu and Mengyin Bao for acquisition of air quality and population data is gratefully acknowledged.

Author Contributions

Y.-L.Z. designed the study, performed data analysis and wrote the manuscript. F.C. participated in data analysis, reviewed and approved the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhang, Y.-L. and Cao, F. Fine particulate matter (PM_{2.5}) in China at a city level. *Sci. Rep.* **5**, 14884; doi: 10.1038/srep14884 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>